

ERRATUM

Bagcilar Med Bull 2023;8(1):112

DOI: 10.4274/BMB.galenos.2023.e002



DOI: 10.4274/BMB.galenos.2022.2022-07-064 upon request from the author “Data on demographic characteristics [body mass index, gender, age, ASA use, preoperative pain value (VAS0)], analgesic use habits, anesthesia technique chosen by anesthesiologist blind to the study, perioperative hemodynamic parameters, additional opioid during surgery, analgesic or anesthetic need, and perioperative complications (patients who willingly stopped the procedure during the intervention, moaning, hypotension, hypertension, bradycardia, respiratory depression, desaturation and need for mask ventilation) were taken from the records of the individuals with the treatment of PKP between July 01 and 22, 2022 after the approval of the University of Health Sciences Turkey, Bursa High Specialization Training and Research Hospital Ethics Committee (2021-KAEK-25 2021/12-09), clinical trials number NCT05526794 and the patient’s consent.” The expression in the Study Design section has been corrected as “Data on demographic characteristics [body mass index, gender, age, ASA use, preoperative pain value (VAS0)], analgesic use habits, anesthesia technique chosen by anesthesiologist blind to the study, perioperative hemodynamic parameters, additional opioid during surgery, analgesic or anesthetic need, and perioperative complications (patients who willingly stopped the procedure during the intervention, moaning, hypotension, hypertension, bradycardia, respiratory depression, desaturation and need for mask ventilation) were taken from the records of the individuals with the treatment of PKP between 01 January-28 June 2022 after the approval of the University of Health Sciences Turkey, Bursa High Specialization Training and Research Hospital Ethics Committee (2021-KAEK-25 2021/12-09), clinical trials number NCT05526794 and the patient’s consent.”.